

Effect of Integrated Nutrient Management on Flowering, Flower Quality and Flower Yield of *Gaillardia pulchella* Foug.) under Hill Zone of Karnataka

Indhumathi M.¹, Chandrashekar S.Y.*², Srinivasa V.³, Shivaprasad M.⁴ and Girish R.⁵

¹PG Scholar, Department of Floriculture and Landscape Architecture,
College of Horticulture, Mudigere (Karnataka), India.

²Associate Professor and Head, Department of Floriculture and Landscape Architecture,
College of Horticulture, Mudigere (Karnataka), India.

³Professor and Head, Department of Vegetable Science, College of Horticulture, Mudigere (Karnataka), India.

⁴Associate Director of Research and Professor of Agronomy, ZAHRS, Mudigere (Karnataka), India.

⁵Scientist (Plant Protection), Krishi Vigyan Kendra, Mudigere (Karnataka), India.

(Corresponding author: Chandrashekar S.Y.*)

(Received: 19 November 2022; Revised: 05 December 2022; Accepted: 20 December, 2022; Published: 09 January, 2023)

(Published by Research Trend)

ABSTRACT: The integrated nutrient management (INM) focuses at efficient and judicious use of the major sources of plant nutrients. It is done in an integrated manner so as to get maximum economic yield without any deleterious effect which is affecting the physio-chemical and biological properties of the soil. Only few experimental study has been done in *gaillardia* for integrated nutrient management. With this background an experiment was conducted at the Experimental block located at the department of Floriculture and Landscape Architecture, College of Horticulture, Mudigere during 2021-22. The experiment was laid out in Randomized Block Design (RBD) having twelve treatments viz., T₁-75% Recommended Dose of Fertilizers + vermicompost @ 1.25 t/ha T₂-50% RDF + vermicompost @ 2.5 t/ha, T₃-75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha, T₄-50% RDF+ vermicompost@ 2.5 t/ha + *Azospirillum* @ 2 kg/ha, T₅-75% RDF + *Azospirillum* @ 2 kg/ha + VAM@2 kg/ha, T₆ - 50% RDF + *Azospirillum* @ 2 kg/ha + VAM @2 kg/ha, T₇-75% RDF+ vermicompost @ 1.25 t/ha + VAM @2 kg/ha, T₈-50% RDF+ vermicompost @ 2.5 t /ha + VAM @2 kg/ha, T₉-75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @2 kg/ha, T₁₀-50% RDF + vermicompost @ 2.5 t/ha + *Azospirillum* @ 2 kg/ha + VAM @2 kg/ha, T₁₁-100% RDF + vermicompost @1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha and T₁₂-100% RDF (Control) replicated thrice. The treatment T₉-75% RDF+ vermicompost@ 1.25 t/ha + *Azospirillum*@ 2 kg/ha + VAM @2 kg/ha recorded the minimum days for 50 per cent flowering (78.30), maximum flowering duration (141.67 days), flower head diameter (6.58 cm), flower petiole length (39.20 cm), corolla length (3.09 cm), shelf life (3.13 days), vase life (5.67 days), average flower weight (3.48 g), number of flowers per plant (104.30), flower yield per hectare (22.09 tonnes)and it was statistically on par with the treatment comprising of T₁₁-100% RDF + vermicompost@ 1.25 t/ha + *Azospirillum*@ 2 kg/ha + VAM@ 2 kg/ha. Hence, this treatment proved to be promising for the enhancement the flower quality and yield of *gaillardia* under hill zone of Karnataka.

Keywords: *Gaillardia*, integrated nutrient management, flowering, flower quality and yield.

INTRODUCTION

Gaillardia (Gaillardia pulchella), a member of the family Asteraceae, is native of Central and Western United States. The generic name *Gaillardia* stands in honour of M. Gaillard, a French patron of botany (Bailey, 1929). It is popularly known as blanket flower. Because of its wide array of colours and patterns including Mexican blankets, gold tipped with russet - red centers but recent introductions have expanded the colour range further. *Gaillardia pulchella* is diploid (2n=36) as reported by Moringa *et al.* (1929). The cultivars of *G. pulchella* have single or double flowers with different colours red, orange yellow, yellow-tipped red, red tipped yellow, bronzed - scarlet and others. Despite its utility for landscape, *gaillardia* is also useful in reduction of soil

erosion in coastal areas (Carig, 1977). *Gaillardia* flourishes well in all kind of garden soils and it can also tolerate high level of light intensities, drought as well as high temperature in a better way as compared to most of the flowering plants. Up to a certain extent it can also tolerant to soil salinity (Tija and Rose 1988).

This is a flower crop which can be utilized as a substitute for chrysanthemum, China aster and marigold etc. of other Asteraceae family crops (Bose *et al.*, 2003). Panchaude (1990) reported that *gaillardia* also has some nematicidal property, it is grown as a catch crop as well as a crop for green manure. The plant has anti-tumour activity owing to the presence of methyl caffeate distributed throughout the plant (Srivastava and Kandpal 2006).

In Karnataka, cultivation of this crop is concentrated in districts such as Belgaum, Dharwad and Bijapur. The growing popularity of this crop has led to expand its cultivation to other parts of the state. The farmers in southern Karnataka are lured by the flower and are braced to take up cultivation of the same on a large scale. In North Karnataka, flowers are used for making *veni* and garlands (Arulmani *et al.*, 2016).

In modern agriculture, the use of chemical fertilizers is an essential. To keep the sustainability of yield it is important, but it poses problems to soil health in long span of time. Therefore, there is a need to restrict the use of chemical fertilizers to certain limit (Gotmare *et al.*, 2007). The integrated nutrient management (INM) focuses at efficient and judicious use of all the major sources of plant nutrients. Nutrients applied in an integrated manner to get maximum economic yield without any deleterious effect which is affecting the physio-chemical and biological properties of the soil.

The nutritional management of gaillardia crop can manifest the performance of the crop. There is an incessant demand for integrated nutrient management. The integrated nutrient management focuses at getting higher yield, quality, and safeguards the interests of farmers and environmental concerns. Hence, the present experimental study was conducted to evaluate the effect of integrated nutrient management on flowering, flower quality and yield of gaillardia.

The multifarious effects of vermicompost influence the growth and yield of crops. *Azospirillum* is a bacterial associative symbiotic biofertilizer contributes in fixation of nitrogen. The vesicular arbuscular mycorrhiza (VAM) is a beneficial fungus that plays an important role in soil nutrient dynamics improving soil physical, chemical and biological properties. VAM is able to alter mobilization of soil phosphorous dynamics in the rhizosphere and mycorrhizal mechanisms (Sharma, 2006).

MATERIALS AND METHODS

Studies were carried out at the experimental block, Department of Floriculture and Landscape Architecture

located at College of Horticulture, Mudigere, Keladi Shivappa Nayaka University of Agriculture and Horticultural Sciences, Shivamogga during 2021 - 22. The soil was prepared to fine tilth. The plot size of 2 m × 1.5 m were prepared under open field condition. Gaillardia seedlings were transplanted at a spacing of 45 cm × 30 cm. The experiment was laid out in randomized block design (RBD) with twelve treatment combinations consisting of inorganic fertilizers (150:80:60 kg NPK), FYM, Vermicompost and bio-fertilizers like VAM, *Azospirillum* with three replications. The different organic manures like FYM, vermicompost and biofertilizers like VAM, *Azospirillum* were supplemented as basal dose according to the treatment to respective plots before transplanting gaillardia seedlings. The recommended dose of chemical fertilizers for gaillardia at 150:80:60 kg NPK per hectare has been applied according to the treatments. 50 per cent dose of nitrogen and full dose of phosphorus and potash were applied as basal dose before transplanting of gaillardia. The balance of 50 per cent dose of nitrogen was applied as top dressing at 30 days after transplanting of the seedlings. Treatments included T₁ - 75 % Recommended Dose of Fertilizers + vermicompost @ 2.5 t/ha, T₂ - 50 % RDF + vermicompost @ 1.25 t/ha, T₃ - 75 % RDF + vermicompost @ 2.5 t/ha + *Azospirillum* @ 2 kg/ha, T₄ - 50 % RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha, T₅ - 75 % RDF + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha, T₆ - 50 % RDF + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha, T₇ - 75 % RDF + vermicompost @ 2.5 t/ha + VAM @ 2 kg/ha, T₈ - 50 % RDF + vermicompost @ 1.25 t/ha + VAM @ 2 kg/ha, T₉ - 75 % RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha, T₁₀ - 50 % RDF + vermicompost @ 2.5 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha, T₁₁ - 100 % RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha and T₁₂ - 100 % RDF (Control). The observations were recorded at 30, 60 and 90 DAT and were statistically analysed.




Fig 1. Effect of integrated nutrient management on number of flowers per plant of gaillardia.

Table 1: Effect of integrated nutrient management on flowering parameters of gaillardia.

Tr. No.	Treatment details	Days taken for visible flower bud formation	Days taken for flower stalk emergence	Days taken for blooming	Days taken for 50% flowering	Flowering duration (days)
T ₁	75 % RDF + VC @ 1.25 t/ha	66.77	68.63	74.63	99.30	123.33
T ₂	50 % RDF + VC @ 2.5 t/ha	67.90	70.77	76.77	102.10	120.67
T ₃	75% RDF + VC @ 1.25 t/ha + Azo @ 2 kg/ha	56.50	59.50	66.20	87.17	135.00
T ₄	50% RDF + VC @ 2.5 t/ha + Azo @ 2 kg/ha	58.43	61.97	68.43	90.43	133.33
T ₅	75% RDF + Azo @ 2 kg/ha + VAM @ 2 kg/ha	69.30	72.23	78.57	103.57	118.67
T ₆	50% RDF + Azo @ 2 kg/ha + VAM @ 2 kg/ha	70.20	74.97	79.97	105.30	116.00
T ₇	75% RDF + VC @ 1.25 t/ha + VAM @ 2 kg/ha	60.80	63.23	71.03	92.90	131.33
T ₈	50% RDF + VC @ 2.5 t/ha + VAM @ 2 kg/ha	62.80	64.47	72.80	94.80	127.67
T ₉	75% RDF + VC @ 1.25 t/ha+ Azo @ 2 kg/ha + VAM @ 2 kg/ha	49.37	53.03	59.03	78.30	141.67
T ₁₀	50% RDF + VC @ 2.5 t/ha + Azo @ 2 kg/ha + VAM @ 2 kg/ha	53.87	57.53	64.20	85.20	136.67
T ₁₁	100% RDF + VC @ 1.25 t/ha +Azo @ 2 kg/ha + VAM @ 2 kg/ha	50.57	55.07	61.13	80.67	138.33
T ₁₂	100% RDF (Control)	64.30	66.47	62.80	97.80	124.67
S. Em±		0.76	0.78	0.80	0.83	1.17
C.D @ 5%		2.24	2.28	2.35	2.44	3.42

RDF = Recommended Dose of Fertilizers VC = Vermicompost Azo = *Azospirillum* VAM= Vesicular Arbuscular Mycorrhiza

Table 2: Effect of integrated nutrient management on flower quality and yield parameters of gaillardia.

Tr. No.	Treatment details	Flower head diameter (cm)	Average flower weight (g)	Shelf life (days)	Flower yield per plant (g)	Flower yield per plot (kg)	Flower yield per hectare (t)
T ₁	75 % RDF + VC @ 1.25 t/ha	4.91	2.95	2.03	237.21	3.32	16.56
T ₂	50 % RDF + VC @ 2.5 t/ha	4.57	2.88	1.90	219.07	3.07	16.12
T ₃	75% RDF + VC @ 1.25 t/ha + Azo @ 2 kg/ha	5.97	3.24	2.80	311.29	4.36	19.89
T ₄	50% RDF + VC @ 2.5 t/ha + Azo @ 2 kg/ha	5.81	3.17	2.65	296.38	4.15	19.20
T ₅	75% RDF + Azo @ 2 kg/ha + VAM @ 2 kg/ha	4.21	2.81	1.77	205.09	2.87	15.89
T ₆	50% RDF + Azo @ 2 kg/ha + VAM @ 2 kg/ha	3.92	2.72	1.56	187.18	2.62	15.46
T ₇	75% RDF + VC @ 1.25 t/ha + VAM @ 2 kg/ha	5.50	3.13	2.44	283.32	3.97	18.49
T ₈	50% RDF + VC @ 2.5 t/ha + VAM @ 2 kg/ha	5.34	3.07	2.31	267.05	3.74	17.85
T ₉	75% RDF + VC @ 1.25 t/ha+ Azo @ 2 kg/ha + VAM @ 2 kg/ha	6.58	3.48	3.13	363.51	5.09	22.09
T ₁₀	50% RDF + VC @ 2.5 t/ha + Azo @ 2 kg/ha + VAM @ 2 kg/ha	6.08	3.32	2.95	328.19	4.59	20.38
T ₁₁	100% RDF + VC @ 1.25 t/ha +Azo @ 2 kg/ha + VAM @ 2 kg/ha	6.40	3.41	3.04	346.12	4.85	21.01
T ₁₂	100% RDF (Control)	5.17	3.03	2.14	256.77	3.59	17.07
S. Em±		0.07	0.04	0.03	6.79	0.10	0.51
C.D @ 5%		0.21	0.13	0.09	19.91	0.28	1.51

RESULTS AND DISCUSSION

The findings of the study were interpreted and listed in Table 1, 2 and Fig. 1 based on the observations recorded.

Effect of integrated nutrient management on flowering parameters

Days taken for visible flower bud initiation and flower stalk emergence. Treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded significantly minimum days after transplanting for first visible flower bud formation (49.37) and flower stalk emergence (53.03 days) which

was on par with the treatment T₁₁ (100% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) 50.57 days for visible flower bud emergence and flower stalk emergence (55.07 days). The treatment T₆ (50% RDF + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded maximum number of days after transplanting (70.20) for visible flower bud initiation and flower stalk emergence (74.97 days). This might be due to the increased C:N ratio. Which resulted in increased the assimilation of photosynthates. This enhanced the source sink relationship and induced early transformation from vegetative to reproductive phase.

Translocated nutrients from shoots to buds reduced the days taken for flowering in gaillardia. The breakdown of apical dominance influenced by enhanced nutrient availability also impacted the faster bud initiation. Phosphorus is the essential nutrient for flowering in plants. The similar findings were reported in experiments done by Parmar *et al.* (2006) in gaillardia, Patel *et al.* (2015) in gaillardia.

Days taken for blooming and 50 per cent flowering.

The plants treated with the treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded significantly minimum days for blooming (59.03) and 50 per cent flowering (78.30 days) which was on par with the treatment T₁₁ (100% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) for blooming (61.13 days) and 50 per cent flowering (80.67 days). Whereas, the treatment T₆ (50% RDF + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) took maximum days for blooming (79.97) and 50 per cent flowering (105.30 days). Application of VAM might have mobilized the phosphorus to rhizosphere, thus increased the P uptake of roots. This might have reduced the days taken for flowering. Gibberellin produced in vermicompost also resulted in early flowering of gaillardia. The results are in line with the research findings of Patil *et al.* (2020) in chrysanthemum, Prasad *et al.* (2018) in dahlia and Bose *et al.* (2018) in china aster.

Duration of flowering. The plant applied with the treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) increased the flowering duration (141.67 days) of gaillardia when compared to control and other treatments. The optimum dose of NPK is needed for induction of early flowering and maximum duration of flowering. Continuous supply of nutrients due to the increased accumulation of them in plant parts encouraged the long duration of flowering. The similar experimental results also reported by Thumar *et al.* (2013) in marigold, Garge *et al.* (2019) in french marigold.

Effect of integrated nutrient management on flower quality parameters of gaillardia

Flower head diameter. The data enumerated in table 2 on flower head diameter in gaillardia as influenced by different nutritional treatments revealed that the treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded significantly maximum flower head diameter (6.58 cm) which was on par with the treatment T₁₁ (100% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) (6.40 cm). Whereas, the treatment T₆ (50% RDF + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded minimum flower head diameter (3.92 cm). Colonization by mycorrhizal fungi also increases root surface area, thus triggers higher nutrient acquisition. Induced source sink relationship also might have promoted the flower quality parameters in gaillardia. The experimental results are in accordance with the studies of Gadagi *et al.* (2004) in marigold, Patel *et al.* (2015) in gaillardia, Koli and Jayanthi (2018) in marigold.

Shelf life. The treatment T₉ recorded significantly maximum shelf life (3.13 days) which was on par with the treatment T₁₁ (3.04 days). Whereas, the treatment T₆ recorded minimum shelf life (1.56 days). Degeneration of water conducting tissues might have reduced by phytohormonal activities of vermicompost, *Azospirillum* and VAM. Shelf life of gaillardia flowers might have extended by biofertilizers. Higher water retention capacity and lesser desiccation of cells exhibited due to the integration of organic manure and biofertilizers might have enhanced the shelf life and vase life of gaillardia flowers. Similar results observed by Kumar *et al.* (2016) in marigold, Ravindra *et al.* (2013) in china aster and Pandey *et al.* (2018) in chrysanthemum.

Effect of integrated nutrient management on yield parameters of gaillardia

Average flower weight. The treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded significantly maximum average flower weight (3.48 g) which was on par with the treatment T₁₁ (100% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) (3.41 g). Whereas, the treatment T₆ (50% RDF + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) recorded minimum average flower weight (2.72 g). It might be due to the activities of vermicompost and biofertilizers. Increased flower diameter and corolla length resulted in maximum individual flower weight. The experimental results are in accordance with the findings of Mittal *et al.* (2010) in marigold, Koli and Jayanthi (2018) in marigold, Kaushik and Singh (2020) in marigold.

Number of flowers per plant. The treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) produced significantly more number of flowers (104.30). This might be due to the increased vegetative growth, as they have produced more number of branches and thus more number of flower buds are initiated compared to control. The increased plant growth accelerated the production of more number of flowers. The results are in accordance with the findings of Gadagi *et al.* (2004) in gaillardia, and Krushnaiah *et al.* (2018) in aster, Abdul *et al.* (2021) in dahlia and Zeighami *et al.* (2015) in Petunia.

Flower yield. The yield parameters *i.e.*, flower yield per plant (363.51 g), per plot (5.09 kg), per hectare (22.09 tonnes) recorded significantly maximum in the treatment T₉ (75% RDF + vermicompost @ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @ 2 kg/ha) as compared to control and other treatments. It might be due to the increased plant growth because of the application of vermicompost, *Azospirillum* and VAM. Growth promoting substances secreted by biofertilizers resulted in increase in yield. Increase in flower yield per plant resulted in maximum flower yield per plot and per hectare. The similar results were also reported by Rathod *et al.* (2002) in gaillardia, Parmar (2006) in gaillardia, Patel *et al.* (2015) in gaillardia, Thumar *et al.* (2013) in marigold and Prasad *et al.* (2018) in dahlia.

CONCLUSION

From the findings of present investigation, it could be concluded that the integrated application of nutrients by

different sources *i.e.*, inorganic fertilizers, vermicopost, *Azospirillum* and VAM influence the plant growth, flowering, quality and yield of gaillardia. Hence, the treatment 75% RDF + vermicompost@ 1.25 t/ha + *Azospirillum* @ 2 kg/ha + VAM @2 kg/ha may be recommended for the commercial cultivation of gaillardia under hill zone of Karnataka.

FUTURE SCOPE

Future studies can be carried out to study the influence of organic mulches and bio-inoculants in gaillardia and to study the effect of integrated nutrient management with pinching in gaillardia.

Acknowledgment. The authors thank the College of Horticulture, Mudigere, KSNUAHS, Shivamogga, for its facilities.

Conflict of Interest. None.

REFERENCES

- Abdul, R. M., Prasad, V. M., Bahadur, V. and Fatmi, U. (2021). Study on Effect of Pinching and Organic Manures on Growth, Flowering and yield of Dahlia (*Dahlia variabilis* L.) cv. Red Symphony. *Biological Forum – An International Journal*, 13(3a), 325-330.
- Arulmani, N., Chandrashekar, S. Y., Ramesha, Y. and Rashmi, R. (2016). Correlation studies in gaillardia (*Gaillardia pulchella* Foug.) genotypes. *Research in Environment and Life Sciences*, 9(4), 458-46.
- Bailey, L. H. (1929). *The Standard Cyclopedia of Horticulture* The MacMillan Company, New York.
- Bose, B. S. C., Prasad, V. M., Prasad, D. S. H. and Sudha, G. (2018). Effect of integrated nutrient management on growth of the China aster (*Callistephus chinensis* L. Nees) cv. Pit and pot. *Plant Archives*, 18(1), 676-678.
- Bose, T. K., Yadav, L. P., Pal, P. Das, P. and Parthasarthy, V. (2003). Commercial flowers. 2nd Edition. *Naya Udyog*, Kolkata, 2, 852-853.
- Carig, R. M. (1977). Herbaceous plant for coastal dune areas. *Proceedings of the Florida State Horticulture Society*, 90, 108-110.
- Gadagi, R. S., Krishnaraj, P. U., Kulkarni, J. H. and Sa-Tong, M. (2004). The effect of combined *Azospirillum* inoculation and nitrogen fertilizer on plant growth promotion and yield response of the blanket flower *Gaillardia pulchella*. *Scientia Horticulturæ*, 100, 323-332.
- Garge, V. C., Malik, S., Awasthi, M., Singh, S. P., Chaudhary, M. and Kumar, A. (2020). Effect of integrated nutrient management on flower quality of French marigold (*Tagetes patula* L.) cv. Pusa Arpita. *International Journal of Pure and Applied Chemistry*, 21(24), 154-158.
- Gotmare, P.T., Damke, M. M., Gonge, V. S. and Deshmukh, S. (2007). Influence of integrated nutrient management on vegetative growth parameters of marigold (*Tagetes erecta* L.). *Asian Journal of Horticulture*, 2(2), 33-36.
- Kaushik, H. and Singh J. P. (2020). Impact of integrated nutrient management (INM) on plant growth and flower yield of African marigold (*Tagetes erecta* L.). *Journal of Pharmacognosy and Phytochemistry*, 9(4), 1481-1484.
- Koli, R. and Jayanthi, R. (2018). Influence of integrated nutrient management on flower yield and economics of marigold (*Tagetes erecta* L.) cv. Pusa Basanti Gainda. *International Journal of Chemical Studies*, 6(4), 2651-2653.
- Krushnaiah, R., Nayak, M. H., Prasanth, P. and Saidanaik, D. (2018). Studies on the effect of integrated nutrient management on growth, flowering and yield of Italian aster (*Aster amellus* L.) cv. 'Purple Multipetal'. *International Journal of Current Microbiology and Applied Sciences*, 7(10), 1-13.
- Kumar, P., Kumar, V. and Kumar, D. (2016). Response of INM to plant growth, flower yield and shelf life of African marigold (*Tagetes erecta* L.) cv. Pusa Basanti Gainda. *International Journal of Agricultural Invention*, 1(1), 108-112.
- Mittal, R., Patel, H. C., Nayee, D. D. and Sitapara, H. H. (2010). Effect of integrated nutrient management on growth and yield of African marigold (*Tagetes erecta* L.) cv. 'Local' under middle Gujarat agro-climatic conditions. *Asian Journal of Horticulture*, 5(2), 347-349.
- Moringa, Y., Fukushima, E., Kano, T. and Yamasaki, Y. (1929). *The Botanical Magazine*, Tokyo, 43, 589.
- Panchaude, M. E. (1990). Proprieties nematocides de quelques plates. *P.H.M. Revue Horticole*, 309, 29-31.
- Pandey, G. Kumar, R., Kumar, S. and Kumar, A. (2018). Effect of integrated nutrient management on floral parameters and soil nutrient status in chrysanthemum (*Chrysanthemum morifolium* Ramat.). *International Journal of Current Microbiology and Applied Sciences*, 7(5), 1984-1990.
- Parmar, V. R. (2006). Influence of bio-fertilizers and nitrogenous fertilizers on growth and flower yield of gaillardia (*Gaillardia pulchella* Foug.) cv. 'Local'. *M. Sc., (Agri.) Thesis* submitted to AAU, Anand.
- Patel, A. S., Leua, H. N., Parekh, N.S. and Patel, H. C. (2015). Effect of integrated nitrogen management on growth, flowering and flower yield of gaillardia (*Gaillardia pulchella* Foug.) cv. 'Lorenziana' under middle Gujarat conditions. *Asian Journal of Horticulture*, 10(1), 126-129.
- Patil, D., Dalal, S. R. and Mahadik, M. K. (2020). Growth, flowering and economics of chrysanthemum cultivation as influenced by integrated nutrient management. *International Journal of Chemical Studies*, 9(1), 313-316.
- Prasad, D. H. S., Prasad, V. M., Goutham, S. K. and Bose, S. C. (2018). Effect of integrated nutrient management on flowering and flower yield of dahlia (*Dahlia variabilis* L.) cv. Kenya Orange. *Plant Archives*, 18(1), 795-798.
- Rathod, N. G., Narwadkar, P. R., Sajindranath, A. K. and Prabu, T. (2002). Effect of integrated nutrient management on growth and yield of gaillardia. *Journal of Maharashtra Agricultural University*, 27(3), 318-319.
- Ravindra, S. P., Hanumanthappa, M., Hegde, J. N., Maheshwar, K. J. and Nagesha, L., (2013). Effect of integrated nutrient management on growth, yield and vase life of China aster (*Callistephus chinensis* L. Nees) for Coastal Karnataka. *Environment and Ecology*, 31(2C), 1104-1106.
- Sharma, A. K. (2006). Biofertilizers for sustainable agriculture, 1st edition. *Agrobios (India)*, Jodhpur. pp. 47-103.
- Srivastava, R. and Kandpal, K. (2006). Gaillardia. In: S. K Bhattacharjee (ed), advances in ornamental horticulture Vol. I, *Pointer Publisher*, Jaipur, pp. 294-304.
- Thumar, B. V., Barad, A. V., Neelima, P. and Bhosale, N. (2013). Effect of integrated system of plant nutrition management on growth, yield and flower quality of African marigold (*Tagetes erecta* L.) cv. Pusa Narangi. *Asian Journal of Horticulture*, 8(2), 466-469.
- Tija, B. and Rose, S. A. (1988). Salt tolerant bedding plants. *Proceedings of Florida State Horticultural Society*, 100, 181-182.
- Zeighami, M., Asgharzadeh, A. and Dadar, A. (2015). Effects of the Biofertilizers Vermicompost and Azotobacter on Qualitative and Quantitative Characteristics of *Petunia hybrid*. *Biological Forum – An International Journal*, 7(1), 586-592.

How to cite this article: Indhumathi M., Chandrashekar S.Y., Srinivasa V., Shivaprasad M. and Girish R. (2023). Effect of Integrated Nutrient Management on Flowering, Flower Quality and Flower Yield of Gaillardia (*Gaillardia pulchella* Foug.) under Hill Zone of Karnataka. *Biological Forum – An International Journal*, 15(1): 119-123.