

## Field Evaluation of Thiamethoxam against Mustard Aphids in Mustard Crop under North Bihar conditions

B. Sujatha<sup>1\*</sup>, Sanjay Kumar Sahoo<sup>2</sup> and Gouri Shankar Giri<sup>3</sup>

<sup>1</sup>Ph.D. Scholar, Department of Entomology, Post Graduate College of Agriculture, Dr. Rajendra Prasad Central Agricultural University, Pusa (Bihar), India.

<sup>2</sup>Associate Professor, Department of Entomology, Post Graduate College of Agriculture, Dr. Rajendra Prasad Central Agricultural University, Pusa (Bihar), India.

<sup>3</sup>Assistant Professor, Tirhut College of Agriculture, Dholi, Dr. Rajendra Prasad Central Agricultural University, Pusa (Bihar), India.

(Corresponding author: B. Sujatha\*)

(Received: 27 November 2022; Revised: 04 January 2023; Accepted: 10 January, 2023; Published: 17 January, 2023)

(Published by Research Trend)

**ABSTRACT:** A field trial was conducted at RPCAU, Pusa, Samastipur, Bihar during the Rabi, 2021-22. Seven treatments were assessed over mustard aphid, *Lipaphis erysimi* Kaltendbach i.e. thiamethoxam 25% WG at different doses at 12.5, 16.67, 25, 37.5, 50 g a.i. per ha, standard check of dimethoate 30% EC at 200 g a.i. per ha and untreated control (water spray). Among the various doses of thiamethoxam sprayed twice at an interval of 10 days on mustard, *Brassica juncea* L., treatments T<sub>5</sub>, T<sub>4</sub> and T<sub>3</sub> were statistically at par in terms of their effectiveness i.e. 5.95, 6.82 and 8.95 no. of *L. erysimi*/top 10 cm of terminal shoot. It was followed by T<sub>6</sub> (13.29), T<sub>2</sub> (15.41), and T<sub>1</sub> (18.75). The plots treated with thiamethoxam (50 g a.i. per ha) gave highest yield and B:C ratio and it was in line with the results of thiamethoxam 37.5 and 25 g a.i. per ha.

**Keywords:** Thiamethoxam, mustard, mustard aphids, field bioefficacy.

### INTRODUCTION

Oil seed crops play an imperative function in Indian agricultural-based economy. Mustard, *Brassica* sp. (*Cruciferae*), farmed as the Rabi crop, is one of the oilseed crops farmed in India (Singh *et al.*, 2018) and major contributor accounting 13 per cent of total consumption of oil worldwide (Giri, 2017). In spite of a diversity of agro-climatic circumstances, along with irrigated/rainfed, timely/late planted, saline soils, and composite farming, it is grown across the country, from the north-eastern/north-western hills to the down south (Pradhan *et al.*, 2020). In India, it is grown on 6.7 million hectares of land, with production of 11.75 million tonnes and productivity of 1524 kg per ha. It is farmed in Bihar on an expanse of 0.08 million hectares, generating 0.10 million tonnes and obtaining yield of 1271 kg per hectare, respectively (Anonymous, 2022).

Mustard is used for various purposes *viz.*, oil, leafy vegetables and seedcake. The leaves contain calcium, copper as well as vitamins- A, C, K and can be eaten in raw or cooked form while their seeds are rich in fiber, selenium, magnesium and manganese. Both seeds

and leaves are rich in sinigrin (anti-inflammatory, antibacterial, antifungal, anticancer and wound-healing properties) and isothiocyanates. Mustard is also rich in carotenoids, isorhamnetin, and kaempferol.

From seedlings to harvests, almost 50 insect species in India pose a serious menace to mustard (Sharma and Singh 2010). Out of many insect pests, sawfly (*Athalia lugens proxima* Klug.), leaf miner (*Chromatomyia horticola* Gorreau), painted bug (*Bagrada cruciferarum* Kirk.), flea beetle (*Phyllotreta cruciferae* Goeze), diamond back moth (*Plutella xylostella* L.), cabbage butterfly (*Pieris brassicae* L.), mustard aphid (*Lipaphis erysimi* Kalt.), cabbage aphid (*Brevicoryne brassicae* L.) and green peach aphid (*Myzus persicae* Sulzer) are considered important causing economic yield losses (Patel *et al.*, 2019). In all mustard-growing countries, the predominant pest is *L. erysimi*. The aphid's nymphs and adults suck cell sap from leaves, stems, twigs, buds, inflorescences, and immature pods, which has a highly deleterious effect on pod setting and production. Nevertheless, aphids secrete honeydew which further causes sooty mould growth, which turns the foliage and pod's appearance filthy black and eventually hampers photosynthetic activity (Awasthi, 2002). For escaping

from the aphid infestation in the mustard crop, there should be asynchronization of the vulnerable stages of the crop and the period of multiplication activity of aphids (Saljoqi *et al.*, 2011). Infestations of aphids peaked from the end of December to the first week in March, according to Patel *et al.* (2019). Hence, if mustard crops are planted early, an infestation can be avoided (Mishra and Gaurav 2022; Saxena and Murthy 2014).

Aphid infestation might diminish oil content by 5 to 6 per cent (Shylesha *et al.*, 2006) and potentially trigger economic output losses amounting to 96 per cent (Patel *et al.*, 2019). The *L. erysimi* is acknowledged as a nationwide pest attributable to its economic importance (Rao *et al.*, 2014). Verma (2000) documented a yield loss of 96 per cent; Bakhetia and Sekhon (1989) observed yield losses as 11.6 to 39.0 per cent; Chauhan and Chauhan (2005) found a loss of 14.0 to 27.9 per cent; and Kular and Kumar (2011) stated a loss of 6.5 to 26.4 per cent; Mukherjee and Singh (2017) estimated the yield losses > 30 per cent. However, Sharma *et al.* (2019) study concludes that for all *Brassica* spp. the mean oil content on late sowing was considerably lower in unprotected plots *i.e.* 38.1 to 38.6 per cent than in protected plots 40.59 to 41.48 per cent.

Insecticides from a more recent generation have lower toxicity toward non-target species, stronger efficacy against the pests they are intended to control, and are not as tenacious as earlier insecticides. Thiamethoxam, a neonicotinoid, is widely used against sucking pests in various crops including mustard (Giri *et al.*, 2017). The efficacy of thiamethoxam in combating mustard aphids is asserted by its broad-spectrum, systemic nature and may be supported by the results of Dhillon *et al.* (2022); Sharma *et al.* (2020); Lal *et al.* (2018); Maurya *et al.* (2018); Shankarganesh *et al.* (2015); Kumar *et al.* (2013). Chemical management is the most effective strategy since the mustard aphid multiplies and spreads quickly in a short amount of time under favourable climatic circumstances. In light of this, the current interpretation was employed to analyze thiamethoxam's field evaluation against mustard aphids in mustard crop under North Bihar conditions.

## MATERIALS AND METHODS

A field experiment was conducted at RPCAU, Pusa, Bihar, India in a Randomized Block Design (RBD) to study the thiamethoxam's field evaluation against aphids, *Lipaphis erysimi* in mustard crop under North Bihar conditions during *Rabi*, 2021-22 with seven treatments *viz.*, T<sub>1</sub>) Thiamethoxam 25% WG @ 12.50 g a.i. per ha; T<sub>2</sub>) Thiamethoxam 25% WG @ 16.67 g a.i. per ha; T<sub>3</sub>) Thiamethoxam 25% WG @ 25 g a.i. per ha; T<sub>4</sub>) Thiamethoxam 25% WG @ 37.50 g a.i. per ha; T<sub>5</sub>) Thiamethoxam 25% WG @ 50 g a.i. per ha; T<sub>6</sub>) Dimethoate 30 % EC @ 200 g a.i per ha (Check); T<sub>7</sub>) Control (water spray) is having each treatment area of 27m<sup>2</sup> with three replications. Sowing of the mustard

crop, Rajendra Sufalam variety was done in October, 2021 according to the standard recommended agronomic practices. Spray solution was calculated with 500 litre of water for one spray for one hectare and in total, two sprays were given with a gap of 10 days. The first application was given when the pest population reached at Economic Threshold Level (ETL). Spraying was done by using knapsack sprayer.

For identification of the mustard aphids, five plants were chosen randomly and tagged. The population of nymphs and adults of aphids were counted from the top 10 centimeters central twig of those plants that were pre-selected. The sightings were identified one day after the first spray (pre-count) as well as after one, three, seven and ten days following every spray. After threshing and sorting the mustard seeds from each plot, the yields were calculated. Per plot's yield was weighted independently and converted to kilograms per hectare then it was analyzed statistically. To combat mustard aphids, the benefit-cost ratio (B: C) of several treatments was computed.

The data on the mustard aphid population in different treatments were subjected to Analysis of Variance (ANOVA) following Randomized Block Design (RBD) using the statistical software OPSTAT (Sheoran *et al.*, 1998). The level of significance was set at 5%.

## RESULTS AND DISCUSSION

The incidence of mustard aphid, before and after two sprays of insecticidal treatments in 2021-22 are illustrated in Table 1. The nymphs and adults aphid population prior to spraying was 92.53-102.93 per top 10 cm terminal shoot. After the first insecticidal application, aphids population was significantly reduced in all the treated plots, but augmented in control plots. Three days after 1<sup>st</sup> application of insecticides spray, results showed that the thiamethoxam (50 g a.i. per ha) treated plot had the least *L. erysimi* (8.47 aphids) followed by thiamethoxam at 37.5 g a.i. per ha (11.20) and thiamethoxam at 25 g a.i. per ha (12.90). Comparatively less effective treatments were dimethoate at 200 g a.i. per ha (18.73) followed by thiamethoxam at 16.67 g a.i. per ha (25.93) and thiamethoxam at 12.5 g a.i. per ha (21.41). Seven days after 1<sup>st</sup> spray application, again least *L. erysimi* plot was thiamethoxam 50, 37.5 and 25 g a.i. per ha (1.95, 2.20 and 5.53). Furthermore followed by dimethoate at 200 g a.i. per ha (9.73 aphids), thiamethoxam at 16.67 g a.i. per ha (13.20) and thiamethoxam at 12.5 g a.i. per ha (17.80). After ten days of 1<sup>st</sup> spray treatment, the population of aphids was increased in all the treatments except in contrast the higher doses of thiamethoxam at 37.5 and 50 g a.i. per ha where the population was reduced. Hence, there was a need to go for the second spray application of the same insecticides.

After the 2<sup>nd</sup> spraying was done, one day after 2<sup>nd</sup> spray, it was noticed that the aphid population was least in

thiamethoxam at 50 and 37.5 g a.i per ha i.e. 0.33 and 0.80 followed by thiamethoxam at 25g a.i per ha (2.40) which was statistically at par. Comparatively less effective treatments were dimethoate at 200 g a.i. per ha (8.52) followed by thiamethoxam at 16.67 g a.i per ha (7.21) and 12.5 g a.i per ha (10.41). Three days after 2<sup>nd</sup> spray appliance, it was reflected in line with the one-day post-application in terms of efficacy, again thiamethoxam at 50 g a.i per ha (0.00) showed a significant reduction in aphid population followed by thiamethoxam at 37.5 g a.i per ha (0.53) and 25g a.i per ha (1.57). Again the comparatively less effective treatments were dimethoate at 200 g a.i. per ha (3.08 aphids) followed by thiamethoxam at 16.67 g a.i per ha (5.71) and 12.5 g a.i per ha (6.13). Seven and ten days after the 2<sup>nd</sup> spray post-appliance, the same trend was followed. Hence the order of efficacy of these treatments was T<sub>3</sub> – thiamethoxam 25 WG @ 50 g a.i. per ha > T<sub>4</sub>-thiamethoxam 25 WG @ 37.5 g a.i. per ha > T<sub>5</sub>-thiamethoxam 25 WG @ 25 g a.i. per ha > T<sub>6</sub> - dimethoate 30 EC @ 200 g a.i. per ha > T<sub>2</sub>-thiamethoxam 25 WG @ 16.67 g a.i. per ha > T<sub>1</sub> - thiamethoxam 25 WG @ 12.5 g a.i. per ha.

The findings of Kumar *et al.* (2022) were comparable to the current observation, which suggested that thiamethoxam 25 WG @ 100 g per ha gave 94.9% reduction and was effective in the treatment of aphids and thiamethoxam 25WG @ 50 g per ha resulted in

71.3% reduction. Din *et al.* (2022) work also supported and they too recorded the lowest no. of aphids in plots treated with thiamethoxam 25 WG @ 100 g per ha in both mustard cultivars *i.e* China (8.49) and Swabi (19.72). The current outcomes were also supported by Kumar (2021) who stated that thiamethoxam 25WG @ 0.2g/ liter water (4.8 aphids/ 10cm central twig) and dimethoate 30 EC @ 1ml/ liter of water (23.6 aphids/ 10 cm central twig) were effective in controlling *L.erysimi*. Additionally, the outcomes of this investigation are in line with Raju and Tayde (2022) who on spraying thiamethoxam 25% WG @ 0.25 g/ lit recorded 33.63 aphids over control (171.96). The conclusions were drawn by Sharma *et al.* (2020); Patel *et al.* (2020); Vishal *et al.* (2019); Dwivedi and Singh (2019); Maurya *et al.* (2018); Lal *et al.* (2018); Vishvendra *et al.* (2018); Kumar *et al.* (2018); Patel *et al.* (2017); Kumar *et al.* (2017); Singh *et al.* (2017); Sen *et al.* (2017) aligned with the current outcomes which say that thiamethoxam 25 WG was most effective in controlling of mustard aphids as well as they have reported the efficacy of dimethoate over aphids.

The data on yield (Kg/ha) is presented in (Table 1) implied that post insecticidal application and the yields were varied significantly from 1172.84 to 1470.74 Kg/ha. The lowest seed yield was reported in control plot yields of 1001.80 Kg/ ha.

**Table 1: Efficacy of insecticides against mustard aphids, Rabi 2021-22.**

Treatments	Number of nymphs and adults of aphids/ top 10 cm shoot											Yield (Kg/ha)	B: C ratio	
	1 <sup>st</sup> spray						2 <sup>nd</sup> spray							Overall Mean
	Pre count	1 DAS	3 DAS	7 DAS	10 DAS	Mean	1 DAS	3 DAS	7 DAS	10 DAS	Mean			
T <sub>1</sub> - Thiamethoxam 25% WG @ 12.50 g a.i/ ha	95.40 <sup>a</sup> (9.82)	56.67 <sup>c</sup> (7.59)	25.93 <sup>c</sup> (5.19)	17.80 <sup>d</sup> (4.34)	23.67 <sup>d</sup> (4.97)	31.02 <sup>d</sup> (5.66)	10.41 <sup>c</sup> (3.38)	6.13 <sup>b</sup> (2.67)	5.14 <sup>b</sup> (2.48)	4.27 <sup>b</sup> (2.30)	6.49 <sup>b</sup> (2.74)	18.75 <sup>c</sup> (4.44)	1172.84	3.384:1
T <sub>2</sub> - Thiamethoxam 25% WG @ 16.67 g a.i/ ha	95.73 <sup>a</sup> (9.84)	49.80 <sup>d</sup> (7.13)	21.41 <sup>bc</sup> (4.73)	13.20 <sup>cd</sup> (3.77)	17.07 <sup>c</sup> (4.25)	25.37 <sup>c</sup> (5.14)	7.21 <sup>bc</sup> (2.87)	5.71 <sup>b</sup> (2.59)	4.96 <sup>b</sup> (2.44)	3.89 <sup>b</sup> (2.21)	5.44 <sup>b</sup> (2.54)	15.41 <sup>bc</sup> (4.05)	1219.75	4.513:1
T <sub>3</sub> - Thiamethoxam 25% WG @ 25 g a.i/ ha	94.40 <sup>a</sup> (9.77)	42.60 <sup>bc</sup> (6.60)	12.90 <sup>a</sup> (3.73)	5.53 <sup>ab</sup> (2.56)	6.39 <sup>b</sup> (2.72)	16.86 <sup>b</sup> (4.23)	2.40 <sup>ab</sup> (1.84)	1.57 <sup>a</sup> (1.60)	0.22 <sup>a</sup> (1.11)	0.00 <sup>a</sup> (1.00)	1.05 <sup>a</sup> (1.43)	8.95 <sup>a</sup> (3.15)	1405.06	8.946:1
T <sub>4</sub> - Thiamethoxam 25% WG @ 37.50 g a.i/ ha	96.07 <sup>a</sup> (9.85)	38.07 <sup>ab</sup> (6.25)	11.20 <sup>a</sup> (3.49)	2.20 <sup>a</sup> (1.79)	1.74 <sup>ab</sup> (1.65)	13.30 <sup>ab</sup> (3.78)	0.80 <sup>a</sup> (1.34)	0.53 <sup>a</sup> (1.24)	0.00 <sup>a</sup> (1.00)	0.00 <sup>a</sup> (1.00)	0.33 <sup>a</sup> (1.15)	6.82 <sup>a</sup> (2.80)	1432.04	9.228:1
T <sub>5</sub> - Thiamethoxam 25% WG @ 50 g a.i/ ha	99.40 <sup>a</sup> (10.02)	36.40 <sup>a</sup> (6.12)	8.47 <sup>a</sup> (3.08)	1.95 <sup>a</sup> (1.72)	0.47 <sup>a</sup> (1.21)	11.82 <sup>a</sup> (3.58)	0.33 <sup>a</sup> (1.15)	0.00 <sup>a</sup> (1.00)	0.00 <sup>a</sup> (1.00)	0.00 <sup>a</sup> (1.00)	0.08 <sup>a</sup> (1.04)	5.95 <sup>a</sup> (2.64)	1470.74	9.757:1
T <sub>6</sub> -Dimethoate 30% EC @ 200 g a.i/ ha	102.93 <sup>a</sup> (10.19)	45.53 <sup>cd</sup> (6.82)	18.73 <sup>b</sup> (4.44)	9.73 <sup>bc</sup> (3.28)	16.20 <sup>c</sup> (4.15)	22.55 <sup>c</sup> (4.85)	8.52 <sup>c</sup> (3.08)	3.08 <sup>ab</sup> (2.02)	2.69 <sup>ab</sup> (1.92)	1.82 <sup>ab</sup> (1.68)	4.03 <sup>ab</sup> (2.24)	13.29 <sup>b</sup> (3.78)	1278.40	4.875:1
T <sub>7</sub> -Control (Water spray)	92.53 <sup>a</sup> (9.67)	98.30 <sup>f</sup> (9.96)	108.93 <sup>d</sup> (10.48)	130.53 <sup>e</sup> (11.47)	158.53 <sup>e</sup> (12.63)	124.08 <sup>e</sup> (11.18)	163.53 <sup>d</sup> (12.83)	178.87 <sup>e</sup> (13.41)	190.80 <sup>e</sup> (13.85)	205.97 <sup>e</sup> (14.39)	184.79 <sup>e</sup> (13.63)	154.43 <sup>d</sup> (12.47)	1001.80	--
F-test	NS	S	S	S	S	S	S	S	S	S	S	S	--	--
SEm±	3.395	1.846	1.810	1.522	1.733	1.455	1.737	1.082	1.326	1.040	1.256	1.299	--	--
C.D (0.05)	N/A	5.751	5.640	4.742	5.398	4.532	5.411	3.370	4.132	3.240	3.913	4.047	--	--
C.V	6.086	6.092	10.574	10.198	9.376	7.199	10.900	6.694	7.890	6.840	7.531	7.043	--	--

NS- Non significant; S- Significant; DAS- Days After Spray; Figures in parenthesis are square root transformed values

The highest seed yield was observed in thiamethoxam at 50 g a.i. per ha (1470.74 Kg/ ha) followed by thiamethoxam at 37.5 g a.i. per ha (1432.04 Kg/ha) and 25 g a.i. per ha (1405.06 Kg/ ha). Despite the fact, that has proven that the greatest dose of thiamethoxam 25 WG at 50 g a.i per ha, was the best treatment for enhancing commercial output, the statistically equivalent dose of 25 g a.i per ha would represent the optimal dose while considering economic as well as prudent usage.

Consequently, it may be ascertained, thiamethoxam 25 WG at 25 g a.i. per ha for limiting the mustard aphid. Kumar (2021) reported the highest seed yield of mustard with thiamethoxam 25 WG @ 0.2 g per liter (1925 kg per ha). This is in line with Kumar *et al.* (2022) who obtained good yields on treatment with thiamethoxam 25 WG @100 g per ha and 50 g per ha. The present results were supported by Sharma *et al.* (2020) who applied thiamethoxam 25 WG (25 g a.i per ha) and obtained a seed yield of 1370 kg per ha.

Among the various tested insecticides cited in the Table.1 it is observed that thiamethoxam at 50 g a.i. per ha (9.757:1), 37.5 g a.i. per ha (9.228:1) and 25 g a.i. per ha (8.946:1) have good benefits out of the cost incurred followed by dimethoate at 200 g a.i. per ha (4.875:1), thiamethoxam at 37.5 g a.i. per ha (1.4513:1) and 12.5 g a.i. per ha (3.384:1). This was supported by the findings of Patel *et al.* (2020).

## CONCLUSION

On brief account of the field evaluation carried out, to cope with the rapidly multiplying aphid population, the insecticidal application would reduce the populations drastically over the control plots. Keeping this in view, firstly we have to follow the practices that would reduce the insecticidal application, and the fact that early sowing would reduce mustard aphid population should be kept in mind while planting. Under unavoidable circumstances like late harvesting of the earlier crop, labour unavailability and tillage practices, even though it is sown lately go for recommended rate of insecticidal applications to avoid judicious usage. Although the highest yield, economics, and lowest aphid population were encountered in plots treated by thiamethoxam 25 WG @ 50 g a.i. per ha followed by 37.5 g a.i. per ha and 25 g a.i. per ha. But, keeping in view of the economic and judicious usage of the insecticides, thiamethoxam 25 WG @ 25 g a.i. per ha could be employed in obtaining good seed yields as well as reducing aphid populations.

## FUTURE SCOPE

To recommend the correct time *i.e.* after the pest population reaches economic threshold level (ETL), the dosage of the insecticides and avoid haphazard and judicious usage of insecticides to protect the natural enemies in the fields.

**Acknowledgements.** I, kindly acknowledge the Professor & Head, Department of Entomology, Post Graduate College of Agriculture, Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar for providing the necessary facilities during my research and financial support from the SRF fellowship awarded by ICAR.

**Conflict of Interest.** None.

## REFERENCES

- Anonymous (2022). <https://www.indiastat.com>. Ministry of Agriculture & Farmers Welfare, Govt. of India.
- Awasthi, V. B. (2002). Introduction to General and Applied Entomology, Scientific publisher, Jodhpur (India), pp. 266-71.
- Bakhetia, D. R. C. and Sekhon, B. S. (1989). Insect-pests and their management in rapeseed-mustard. *Journal of Oil seeds Research*, 6, 269-299.
- Chauhan, S. and Chauhan, S. V. S. (2005). Quantity and quality of oil in Indian mustard (*Brassica juncea* L. Czern and Coss) grown under various kinds of stress. *Brassica*, 7(1), 59-62.
- Dhillon, M. K., Singh, N. and Yadav, D. K. (2022). Preventable yield losses and management of mustard aphid, *Lipaphis erysimi* (Kaltenbach) in different cultivars of *Brassica juncea* (L.) Czern & Coss. *Crop Protection*, 161, 106070.
- Din, M., Khan, S. A., Shah, S. H. and Ullah, N. (2022). Efficacy of different pesticides against mustard aphid, *Lipaphis erysimi* in mustard cultivars. *Pakistan Journal of Agricultural Research*, 35(2), 371-379.
- Dwivedi, S. A. and Singh, R. S. (2019). Evaluation of new molecular insecticides and biopesticides against mustard aphid, *Lipaphis erysimi* (Kalt.) on yield parameter in mustard. *Journal of Biopesticides*, 12(2), 203-214.
- Giri, G. S. (2017). *Effect of thiamethoxam on growth, development and behavior of Apis mellifera* L. Diss. M. Sc. Thesis. GB Pant University of Agriculture and Technology Pantnagar, Uttarakhand, India, 2017.
- Giri, G. S., Mall, P. and Pandey, R. (2017). Effect of thiamethoxam on colony development of *Apis mellifera* L. *Journal of Entomology and Zoological studies*, 5(6), 177-179.
- Kular, J. S. and Kumar, S. (2011). Quantification of avoidable yield losses in oilseed *Brassica* caused by insect pests. *Journal of Plant Protection Research*, 51(1), 38-43.
- Kumar, H., Singh, S. and Yadav, A. (2022). Bio-efficacy of insecticides against mustard aphid (*Lipaphis erysimi*) in Indian mustard (*Brassica juncea* L.). *Journal of Oilseeds Brassica*, 13(2), 158-162.
- Kumar, K., Sachan, S. K. and Singh, D. V. (2013). Bio-efficacy of some new insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.) and their effect on coccinellid population in rapeseed mustard. *Vegetos-International Journal of Plant Research*, 26(2), 159.
- Kumar, S. (2021). Evaluation of insecticides for their field efficacy against turnip aphid, *Lipaphis erysimi* (Kaltenbach) on Indian mustard. *Journal of Oilseeds Brassica*, 12(2), 113-116.
- Kumar, S. R., Singh, V., Yadav, S. K. and Yadav, A. K. (2017). Efficacy of insecticides and botanicals against *Lipaphis erysimi* (Kalt.) on mustard. *Indian Journal of Entomology*, 79(4), 503-506.
- Kumar, A., Srivastava, A., Rawal, R. and Roy, N. (2018). Chemical evaluation against mustard in Bhundelkhand region. *Journal of Entomology and Zoology studies*, 6(2), 3195-3197.
- Lal, B., Nayak, M. K., Tomar, D. S. and Thakur, S. R. (2018). Efficacy of newer insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.) in Indian mustard under

- Bundelkhand Agro climatic zone of Madhya Pradesh. *Journal of Entomology and Zoology Studies*, 6(6), 400-403.
- Maurya, N. K., Singh, R., Singh, J., Nigam, R., Hasan, W. and Kumar, A. (2018). Efficacy of novel insecticides against mustard aphid *Lipaphis erysimi* (Kaltenbach). *International Journal of Agricultural Invention*, 3(1), 62-70.
- Mishra, M. and Gaurav, K. (2022). Management of mustard aphid (*Lipaphis erysimi* Kalt.) by manipulating sowing dates. *Journal of Pharmaceutical Negative Results*, 13(9), 8489-8498.
- Mukherjee, U. and Singh, P. P. (2017). Assessment of yield losses in rape-seed mustard due to mustard aphid (*Lipaphis erysimi* Kalt.) in Bihar. *Trends in Biosciences*, 10(41), 8655-8659.
- Patel, S., Singh, C. P. and Hasan, W. (2020). Relative efficacy of certain insecticides against mustard aphid in mustard ecosystem. *International Journal of Agricultural and Applied Sciences*, 1(1), 46-48.
- Patel, S., Singh, C. P. and Yadav, S. K. (2019). Monitoring of insect-pest complex on rapeseed-mustard at Pantnagar. *Journal of entomological research*, 43(1), 73-76.
- Patel, S., Yadav, S. K. and Singh, C. P. (2017). Bio-efficacy of insecticides against *Lipaphis erysimi* (Kalt.) in mustard ecosystem. *Journal of Entomology and Zoology studies*, 5(2), 1247-1250.
- Pradhan, P. P., Borkakati, R. N. and Saikia, D. K. (2020). Insect pests of mustard and their natural enemies in Assam. *International Journal of Current Microbiology and Applied Sciences*, 9(7), 2785-2790.
- Raju, C. E. P. and Tayde, A. R. (2022). Field evaluation of selected insecticides and botanical against mustard aphid, *Lipaphis erysimi* (Kalt.) on mustard, *Brassica juncea* L. *International Journal of Plant & Soil Science*, 34(22), 1188-1193.
- Rao, B. B., Rao, V. U. M., Nair, L., Prasad, Y. G., Ramraj, A. P. and Chattopadhyay, C. (2014). Mustard aphid infestation in India development of forewarning models. *Journal of Environmental Biology*, 35, 683-688.
- Saljoqi, A.R., Zada, R. and Khan, I. A. (2011). Population trend of canola aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae) and its associated natural enemies in different *Brassica* lines along with the effect of gamma radiation on their population. *World Acad. Sci., Engin. Tech.*, 73, 187-194.
- Saxena, S. and Murty, N. S. (2014). Weather based model development or outbreak of mustard aphid (*L. erysimi* Kalt.) using artificial neural network. *The Bioscan*, 8(1&2), 47-52.
- Sen, K., Samanta, A., Hansda, A., Dhar, P. P. and Samanta, A. (2017). Bioefficacy and economics of some insecticides against mustard aphid, *Lipaphis erysimi* (Kalt.) infesting mustard. *Journal of Crop and Weed*, 13(2), 235-237.
- Shankarganesh, K., Suroshe, S. S. and Paul, B. (2015). Relative susceptibility of the Bikaner and Delhi populations of mustard aphid, *Lipaphis erysimi* (Kalt.) (Homoptera: Aphididae), and its predator, *Coccinella septempunctata* L. (Coleoptera: Coccinellidae), to different insecticides. *Phytoprotection*, 95(1), 1-47.
- Sharma, D., Yadav, S. P. and Yadav, S. (2019). Assessment of avoidable yield losses due to *Lipaphis erysimi* Kalt. in various *Brassica* spp. *International Journal of Current Microbiology and Applied Sciences*, 8(6), 2020-2025.
- Sharma, N., Upadhyaya, S. N., Singh, U. C., Dubey, M. and Ahmad, A. (2020). Bio efficacy of insecticides against mustard aphid. *Journal of Pharmacognosy and Phytochemistry*, 6, 187-192.
- Sharma, P. and Singh, Y. P. (2010). Directorate of Rapeseed-mustard, Indian Council of Agriculture Research, Sewar, 321303, Bharatpur (Rajasthan). *Annual National Language Journal*, 1, 47-51.
- Sheoran, O. P., Tonk, D. S. Kaushik, L.S., Hasija, R. C. and Pannu, R. S. (1998). Statistical doftware package for agricultural research workers. In: Hooda DS & Hasija RC (eds). Recent advances in information theory, statistics and computer applications, Department of Mathematics and Statistics, CCS HAU, Hisar, pp.139-143.
- Shylesha, A. N., Azad Thakur, N. S., Pathak, K. A., Rao, K. R., Saikia, K., Surose, S., Kodandaram, N. H. and Kalaishekar, A. (2006). Integrated management of insect pest of crops in north eastern hill region. Technical Bulletin No. 19. ICAR RC for NEH Region, Umiam, pp. 50.
- Singh, H., Deshwal, H. L., Haldhar, S. M. and Singh, V. (2017). Bio-efficacy of some insecticides and botanicals against mustard aphid, *Lipaphis erysimi* Kalt. (Hemiptera: Aphididae) on Indian mustard. *Journal of Experimental Zoology, India*, 20(1), 249-253.
- Singh, V. V., Agarwal, N., Satish, B. N., Kumar, S., Sachin, K. and Pal, K. (2018). Studies on insect diversity in mustard (*Brassica campestris* L.) ecosystem. *Journal of Entomology and Zoology Studies*, 6(5), 698-702.
- Verma, K. D. (2000). Economically important aphids and their management. In: IPM System in Agriculture (Eds: Upadhyay, R.K., Mukherji, K.G., Dubey, O.P.). Vol. 7. Aditya Books Pvt. Ltd., New Delhi: pp. 144-162.
- Vishal, Singh, H. and Kumar, A. (2019). Efficacy and economics of some newer insecticides against mustard aphid, *Lipaphis erysimi* (Kalt). *Journal of Pharmacognosy and Phytochemistry*, 8(3), 785-788.
- Vishvendra, Kumar, S., Sachan, S. K., Singh, G. and Singh, R. (2018). Bio-efficacy of insecticides and biorational against *Lipaphis erysimi* (Kalt.) in mustard crop. *Journal of Pharmacognosy and Phytochemistry*, 5, 58-61.

**How to cite this article:** B. Sujatha, Sanjay Kumar Sahoo and Gouri Shankar Giri (2023). Field Evaluation of Thiamethoxam against Mustard Aphids in Mustard Crop under North Bihar conditions. *Biological Forum – An International Journal*, 15(1): 336-340.